





HELMINTHS

FLUKES (TREMATODES)



FLATWORMS

TAPEWORMS (CESTODES)



TAENIA SOLIUM

ROUNDWORMS (NEMATODES)




ENTEROBIUS VERMICULARIS


	TREMATODES	CESTODES	NEMATODES
SHAPE	UNSEGMENTED PLANE	SEGMENTED PLANE	CYLINDRICAL
BODY CAVITY	No	No	Yes
DIGESTIVE TRACT	INCOMPLETE	No	COMPLETE
SEX	HERMAPHRODITES (EXCEPT SCHISTOSOMA)	HERMAPHRODITES	MALE OR FEMALE

HELMINTHS


FLUKES (TREMATODES)




TAPEWORMS (CESTODES)




ROUNDWORMS (NEMATODES)



COMPARATIVE LIFE CYCLE OF INTESTINAL NEMATODES



SOIL - TRANSMITTED HELMINTHS (STHs)



MOST COMMON STHs :

- ASCARIS LUMBRICOIDES
- ANCYLOSTOMA DUODENALE, NECATOR AMERICANUS
- TRICHURIS TRICHIURA

DIAGNOSIS



EXAMINE **STOOLS** OR **TISSUES** FOR **EGGS**, **LARVAE**, OR **ADULT WORMS**

TREATMENT

ANTI PARASITIC MEDICATIONS



Armando H. Faigl